

FACT SHEET

Ankle-Foot Orthoses and Footwear for Children with Cerebral Palsy-Selecting Optimal Designs

An Ankle-Foot Orthosis (AFO) is one that encompasses the ankle joint and the whole or part of the foot.^{1,2} Ankle-Foot Orthoses are worn with footwear, which is integral to biomechanical control, so the overall orthosis is now described as an Ankle-Foot Orthosis Footwear Combination (AFOFC). This factsheet provides advice on the selection of AFOs and footwear used for sitting, standing and walking activities.

This factsheet seeks to provide answers to these questions:

1. How do I begin?
2. How do I incorporate evidence into my clinical decision making?
3. What types of AFOs and footwear are appropriate to consider?
4. How do I determine optimal design and alignments and when the child needs to wear it?

How do I begin?

Orthotic interventions need an interdisciplinary family-centered approach with shared goal setting. Child and family-centered care should include the parents, child, and all professionals involved to share expertise. The key to goal setting and providing optimum interventions is to understand the natural history and prognosis of the condition.

Prognosis is particularly pertinent to orthotic interventions, as the orthosis may provide a preventative role in the natural history of the condition. While the primary insult in Cerebral Palsy (CP) may be static, secondary skeletal and musculotendinous pathologies develop, as the child grows with the effects of the neurological impairments.^{3,4} These may be coupled with the development of pain, an important indicator of quality of life.^{5,6} Cerebral Palsy is a heterogeneous condition, so additional diagnostic information is required to fully understanding prognosis and set appropriate goals; topography, motor function impairment, degree of impairment.⁷⁻¹⁵

How do I incorporate evidence into my clinical decision making?

Evidence-informed practice is the integration of best research evidence with clinical expertise and patient values.^{16,17} Currently, there is limited and low level research evidence on the effectiveness of ankle-foot orthoses for children with CP.¹⁸⁻²⁷ Research has focused on limited aspects of the International Classification of Functioning, Disability and Health (ICF),²⁸ and has been mainly conducted in gait laboratories rather than in daily living activities. Patient values and clinical expertise should be the combined main evidential factors that direct goal setting and clinical decision making. The goals and outcomes for children with CP will span the whole of the ICF.

Reviews¹⁸⁻²⁶ and subsequent studies²⁹⁻³³ report that AFOs can positively influence the arch of the foot (foot posture), gross motor function, spatial temporal measures, kinematics and kinetics, muscle operating lengths and gait efficiency. Ankle power, as calculated by 3D analysis, has been found to be reduced in AFOs that

restrict ankle motion. This is inevitable and is often an acceptable compromise in order to optimize other parameters of gait and functional tasks.²⁰

Increased internal rotation of the feet may be observed, because the effects of the torsions of the long bones are unmasked once pronation is corrected in the AFO.³³ Previous literature, not included in the reviews, has documented positive facilitation of motor learning with optimally aligned AFOFCs.^{34,35} Reviews comment that optimum orthosis and footwear designs and alignments may not have been used in studies.^{20,27} Thus, research must be evaluated in light of that finding.

What types of AFOs and footwear are appropriate to consider?

There are many different designs and types of AFOs. There is now agreed international terminology for categorization and description of orthoses.³⁶ However, many AFO designs were given names previously, and these terms remain in common use.³⁷

Table 1 illustrates frequently used designs and common names used in the literature. Terminology mainly relates to the biomechanical control provided at the ankle joint in the sagittal plane, but all AFO designs and footwear will have sagittal, coronal, transverse and tri-planar design features at ankle and foot joints and all planes need consideration.

How do I determine the optimal design and alignments and when the child needs to wear it?

Determine and Set Goals: First determine the short, medium, and long term goals for the child in collaboration with the family and team. Goals should consider all areas of the ICF (Tables 2-5).³⁸⁻⁴² Once goals are identified, it is possible to determine the AFOFC design, alignments and frequency of use to achieve goals.










Determine optimum design of AFOFC to achieve goals: Tables 2-5 offer design considerations for achieving ICF goals and provide indications and contraindications for commonly used designs of AFOFCs. These tables consider bones and joints, muscles, changing atypical standing and walking patterns, and activities and participation, respectively. Clinical algorithms, based on the indications and contraindications, have been developed, and these may further assist clinical decision making.^{27,38,39,43,44}

Determine optimum alignment of joints and segments in AFOs and AFOFCs: Optimizing joint alignment within an AFO and consideration of the Shank to Vertical Angle (SVA) alignment of the AFOFC is essential. Non-optimal joint and segment alignments will affect sitting, standing, and walking activities, and long term outcomes.^{27,38,44,45} If AFOFCs are used for standing and walking, the design of the heels and soles of the footwear will affect the ability of the AFOFC to change kinematics of segments and joints and kinetics (Table 4).^{27,38,43,44,46}

Determine segment proportion: Optimizing segment proportion usually includes optimizing the length of the foot for height and equalizing any true leg length discrepancy when wearing AFOFCs or footwear.^{38,43}

Determine the dosage required to achieve the goals: Dosage, the percentage of time a child wears the AFOFC to achieve their goals, will need to balance all goals. Some children may need to wear AFOFC for 100% of the week. Other children will have time in the day or week when they do not wear an AFOFC. During these periods, children may still wear footwear with or without a foot orthosis. Design of AFOFCs, of the footwear, and their frequency of use requires regular review.

Table 1. – Designs & Terminology for AFOs, Footwear and AFOFC

AFO DESIGN or TYPE & TERMINOLOGY							
Commonly Used AFO Designs Straps and pads are not illustrated, for clarity.							
Common Name	^a Supra-Malleollar AFO	Articulated or Hinged AFO Articulated AFO with anterior shell not illustrated		Flexible or Spring AFO	^b Solid or Fixed or Rigid AFO ^c AFO with an anterior shell is often called a Ground Reaction or Floor Reaction Orthosis (GRO/ FRO/GRAFO)		
Available motions within orthosis	Ankle: full or partial movement; free, limit, stop, assist or resist designs. ^d Metatarsal phalangeal joints (MTPJs): typically free but may have MTPJs fixed				Ankle: no movement at ankle. MTPJs: free or fixed, GROs -usually MTPJs fixed design.		
FOOTWEAR DESIGNS & TERMINOLOGY - SAGITTAL PLANE							
Heel Sole Pitch	^e Heel Sole Differential (HSD)						
Sole Design	^f Stiffness = Flexible or Stiff ^g Profile = Flat or Rounded ^h Length of Toe Lever				Stiffness = Flexible or Stiff Profile = Flat or Rounded or Point Loading Length of Toe Lever		
Heel Design	Stiffness = Flexible or stiff Profile = Flat or Rocker Length of Heel Lever				Stiffness = Flexible or Stiff Profile = Flat or Rocker Length of Toe Lever		
FOOTWEAR DESIGNS & TERMINOLOGY – CORONAL PLANE							
Sole & Heel Designs	ⁱ Medial and Lateral Flares Stiffness, profile and length						
AFO & AFOFC ALIGNMENTS - SAGITTAL PLANE							
	^j Ankle Angle alignment AFO 				^k Shank to Vertical Angle alignment AFOFC 		

- a. The terms 'Supramalleolar Orthosis' (SMO) and 'Dynamic AFO' (DAFO) emerged in the 1980s. DAFO was usually used to describe an SMO with a neurophysiological footplate, but Dynamic AFO is also used to describe any type of AFO with this design feature.
- b. Fixed Ankle AFO, Rigid Ankle AFO, and Solid Ankle AFO are designs intended to prevent movement at the ankle joint.
- c. Ground Reaction Orthosis (GRO) is an AFO designed to manipulate the Ground Reaction Force (GRF) at the knee. All AFOs have the potential to manipulate the GRF, but 'GRO' is usually linked to an AFO with an anterior shell.
- d. If MTPJs are fixed in an AFO or Footwear and walking requires some 'heel rise', a rocker profile is essential. Design and position of the rocker need to be determined.³⁸
- e. The Heel Sole Differential (HSD) is the difference between the height of the heel of the footwear (rear foot height) and the thickness of the sole at the MTPJs. It is a measure that reflects the pitch of the footwear.³⁸
- f. Stiffness, resistance to bending. MTPJ free designs require a flexible sole; MTPJ fixed require a stiff sole with a rocker profile, rounded or point loading.³⁸
- g. The profile is the shape of the distal surface of the footwear. If sole is stiff, the profile is described as a rocker.³⁸
- h. Heel and Toe Levers (Sagittal Plane); Heel Lever = length of heel of footwear from ankle joint. Toe Lever= length of sole of footwear to MTPJs if flexible and to the rocker position if stiff. The length of heel and toe levers will be determined individually. If the heel lever is long but not stiff the effectiveness as a 'heel lever' is minimal.³⁸
- i. Medial and Lateral Flares (Coronal Plane). The medial and lateral walls of the lowers of the footwear can be extended to produce medial and lateral levers, at the heel or forefoot. They will be of optimum length for the individual and to be effective will be stiff with a flat profile
- j. Ankle Angle alignment of the AFO (AA AFO); determined by length and tone of gastrocnemius, and foot posture.³⁸
- k. Shank to Vertical Angle alignment of the AFOFC (SVA AFOFC) is a measure of the static alignment of the AFOFC when standing still and weight-bearing equally between heel and sole. Optimal SVA AFOFC for an individual child is determined by; gait pattern, degree of hip and knee flexion contractures. Optimal SVAs for standing and walking are usually inclined, most by 10-12° incline, many by 12-15° incline and some by 15-19° incline.³⁸

TABLE 2. DESIGN CONSIDERATIONS FOR BONE & JOINT GOALS



2a

2b

2c

2d

2e

Considerations:

- Abnormal forces imposed on a developing skeleton can lead to abnormal bone growth and joint deformity, predisposing the child to pain.⁴
- Arches of the foot, foot posture, develop during the first 7 years.^{47,48} During this time, feet are susceptible to deformation.
- Within AFOFCs, foot joints and arches need to be in a stable alignment (2a, 2b, 2d) and not deformed (2c). If it is not possible to achieve normal foot posture alignments, optimal alignments of the foot joints are required. In fixed ankle AFOFCs, ankle joint alignment needs to be appropriate for calf muscle length, stiffness, and foot posture;^{27,38,43} in articulated AFOFCs, dorsiflexion should occur at the ankle (2a, 2b), not by foot pronation/supination or deformation of foot joints (2c).^{38,44,49}
- Walking patterns may influence development of bones and joints proximal to the foot and ankle (2e).^{38,50}
- Standing and walking activities require optimum length of foot for height. Children with CP may have a shorter foot for their height and may need the effective foot length optimised in AFOFC prescriptions.³⁸ Leg length discrepancies from short shank or thigh segments usually need to be equalised to prevent abnormal compensatory foot postures and gait patterns from developing.³⁸

AFOFC DESIGN	INDICATIONS	CONTRAINDICATIONS
Ankle: Solid/ Fixed/Rigid	<ul style="list-style-type: none"> • Poor or unstable triplanar alignment of foot • Dorsiflexion occurs at foot joints, not ankle joint • Required range of ankle motion (ROM) not available 	
Ankle Dorsiflexion: Free, Limit, Assist, Resist, Stop	<ul style="list-style-type: none"> • Required ankle joint ROM available • Stable triplanar alignment of the foot • Dorsiflexion occurs at ankle joint, not foot joints 	<ul style="list-style-type: none"> • Required ankle joint ROM not available • Unstable triplanar alignment of the foot • Dorsiflexion occurs at foot joints, not ankle joint
Ankle Plantarflexion: Free, Limit, Assist, Resist, Stop	<ul style="list-style-type: none"> • Required ankle joint ROM available 	<ul style="list-style-type: none"> • Required ankle joint ROM not available
MTPJs: Free	<ul style="list-style-type: none"> • Required MTPJ joint ROM available 	<ul style="list-style-type: none"> • Required MTPJ joint ROM not available
MTPJs: Fixed	<ul style="list-style-type: none"> • MTPJ joint ROM not available • Control of correctable Hallux Valgus • Short foot for height 	

TABLE 3. DESIGN CONSIDERATIONS FOR MUSCLE & TENDON GOALS


					
Muscle actions lower leg				Muscle lengthening	
0-10%GC	10-30% GC	30-50% GC	50-60% GC	40%GC	50% & 100% GC
Considerations: <ul style="list-style-type: none">Goals for muscles within and more proximal to the AFOFC will include muscle length, strength, tone, selective control and timing.^{38,39}Pay attention to tri-articular gastrocnemius, if knee extension is required in standing and walking, as non-optimum ankle angle, may leave insufficient muscle length available for knee extension.^{38,43,44}AFOFCs will need to compensate for weak muscles, have optimal alignments for short and stiff muscles, and induce optimum muscle actions and lengthening when standing and walking.^{38,43,44}					
AFOFC DESIGN		INDICATIONS			
Ankle: Solid/ Fixed/ Rigid		<ul style="list-style-type: none">Weak plantarflexors or dorsiflexorsStiffness/high tone in plantar or dorsiflexorsShort plantarflexors or dorsiflexorsTurn tri-jointed gastrocnemius into uni-jointed muscle to achieve control of lengthening and actions during the gait cycle			
Articulated or Hinged Flexible or Spring		<ul style="list-style-type: none">Indications and contraindications for muscles using these designs are highly complex, so selection requires advanced knowledge of kinematics, kinetics, and muscle actions.			
		INDICATIONS		CONTRAINDICATIONS	
Ankle: Plantarflexion Stop at 90° Dorsiflexion Free		<ul style="list-style-type: none">Weak dorsiflexorsPlantarflexor length: at least 10° dorsiflexion in knee extensionPlantarflexor tone: not marked or considerably increasedPlantarflexor strength: sufficient strength to control shank kinematics & ankle joint dorsiflexion		<ul style="list-style-type: none">Plantarflexor length: short, less than 10° dorsiflexion with knee extended.Plantarflexor tone: marked or considerably increasedPlantarflexor strength: weak, unable to control shank kinematics and ankle joint dorsiflexion	
Ankle: Plantarflexion Free Dorsiflexion Free		<ul style="list-style-type: none">Dorsiflexor strength: Strong in swing phase & 0-10% stance phasePlantarflexor length: at least 10° dorsiflexion with knee extensionPlantarflexor tone: not marked or considerable increasePlantarflexor strength: sufficiently strong to control shank kinematics & ankle joint dorsiflexion		<ul style="list-style-type: none">Dorsiflexor strength: Weak, insufficient to control swing phase & 0-10% stance phasePlantarflexor length: short, less than 10° dorsiflexion with knee extensionPlantarflexor tone: ‘marked or considerable increase’Plantarflexor strength: weak, unable to control shank kinematics and ankle joint dorsiflexion	
MTPJs: Free		Muscle length available		Muscle length not available	
MTPJs: Fixed		Muscle length not available			

TABLE 4. DESIGN CONSIDERATIONS FOR MOTOR CONTROL OF STANDING & WALKING and CHANGING ATYPICAL GAIT PATTERN GOALS



								
Standing & Swaying			Walking - Stepping			Walking - Full Gait Cycles		
Considerations:								
<ul style="list-style-type: none">Children with CP walk with atypical gait patterns for various reasons, including muscle and joint deformity and poor motor control.^{4,27,38} Goals are often set for improving motor control of standing and walking to achieve optimal standing and walking patterns: standing and swaying; stepping, walking with an abbreviated gait cycle, or walking with full gait cycles.^{38,43,44} These patterns may first be achieved when wearing the AFOFC, but later may be achieved using footwear or when barefoot.Motor learning of more typical kinematics and kinetics requires repetitive practice of the activity. Reducing the ‘degrees of freedom’ in the lower limb, by reducing the number of joints that move, may help motor learning. Therefore, fixing the ankle joint in an AFO to learn control of knee and hip may be helpful.^{27,34,35,38}When prescribing AFOFCs for standing and walking activities, consideration of alignment of body segments is as important as the alignment of joints. Normalizing distal segment alignments will often normalise proximal segment and joint alignments, kinematics and kinetics.^{27,38,43,44}								
AFOFC DESIGN		INDICATIONS				CONTRAINDICATIONS		
Ankle: Solid/Fixed/Rigid		<ul style="list-style-type: none">Inability to produce normal kinematics and kinetics of trunk, hip, thigh, knee, shank, foot, if ankle free to moveReduction of ‘degrees of freedom’ from 3 to 2 by fixation of ankle joint; knee and hip joints free to move						
Ankle Dorsiflexion: Free, Limit, Assist, Resist, Stop		<ul style="list-style-type: none">Able to walk with normal segment and joint kinematics and kinetics with movement of ankle joint in selected AFO designNormal foot, shank, knee, thigh, hip, pelvis and trunk kinematics in selected AFOFC				<ul style="list-style-type: none">AFOFC does not correct gait pattern sufficiently.		
Ankle Plantarflexion: Free, Limit, Assist, Resist, Stop								
MTPJs: Free		<ul style="list-style-type: none">Able to normalise walking pattern with MTPJs freeFoot correct length for height				<ul style="list-style-type: none">AFOFC does not correct gait pattern sufficiently		
MTPJs: Fixed		<ul style="list-style-type: none">Foot is short for height; optimal length of foot for height required to achieve normal kinematics and kineticsUnable to normalise walking pattern with MTPJs free.						

TABLE 5. DESIGN CONSIDERATIONS FOR ACTIVITIES & PARTICIPATION GOALS



Considerations:

- A variety of footwear and orthosis for various activities are needed.
- For recreational activities, a general rule, use the orthosis or footwear that allows the best performance of the activity, as long as other ICF goals are not compromised or harm done.
- For ambulant children, there may be complex considerations to balance when deciding the percentage of time an AFOFC is worn.^{38,39}
 - Walking is a common activity and goals for bones, joints, muscles, and development of an optimum walking pattern may depend on the hours of walking in the AFOFC.
 - However, the child will also be undertaking other functional activities in their day, and for many children, there will be periods in the day when they would not use the AFOFC for walking.
 - Often a compromise needs to be made as to when the AFOFC is worn and when not.
- Considerations will be different for differing ages.

AFOFC DESIGN	INDICATIONS	CONTRAINDICATIONS
Ankle: Fixed/Solid/Rigid MTPJs: Free/Fixed	<ul style="list-style-type: none"> • Activity is not possible or performed less well with less support • Goals in other domains of ICF will be compromised in an AFO with less biomechanical control 	
Ankle Dorsiflexion: Free, Limit, Assist, Resist, Stop. Ankle Plantarflexion: Free, Limit, Assist, Resist, Stop	<ul style="list-style-type: none"> • Activity is performed better • Goals in other domains of ICF will not be compromised 	<ul style="list-style-type: none"> • Forces on the muscles and joints would cause harm to developing structures for duration of use in the activity

REFERENCES

1. International Organization for Standardization (ISO) 8549-1. *Prosthetics and orthotics - Vocabulary - Part 1: General Terms for external limb prostheses and external orthoses*; 1989. 10 p.
2. International Organization for Standardization (ISO) 8549-3. *Prosthetics and orthotics - Vocabulary - Part 3: Terms relating to external orthoses*; 1989.
3. Bleck EE. *Orthopaedic Management in Cerebral Palsy*. London: Mac Keith Press; 1987.
4. Gage JR, Schwartz MH, Koop SE, Novacheck T, editors. *The Identification and Treatment of Gait Problems in Cerebral Palsy, 2nd Edition*. London: Mac Keith Press; 2009.
5. Colver A, Rapp M, Eisemann N, Ehlinger V, Thyen U, Dickinson HO, et al. Self-reported quality of life of adolescents with cerebral palsy: a cross-sectional and longitudinal analysis. *Lancet (North American edition)*. 2015; 385(9969):705-16.
6. Parkinson KN, Gibson L, Dickinson HO, Colver AF. Pain in children with cerebral palsy: a cross-sectional multicentre European study. *Acta Paediatr*. 2010; 99(3):446-451.
7. Rosenbaum P, Rosenbloom L. *Cerebral Palsy. From Diagnosis to Adult Life. 1st ed*: Mac Keith Press; 2012. p 224.
8. Sanger TD, Chen D, Delgado MR, Gaebler-Spira D, Hallett M, Mink JW. Definition and classification of negative motor signs in childhood. *Pediatrics*. 2006;118(5):2159-67.
9. Sanger TD, Chen D, Fehlings DL, Hallett M, Lang AE, Mink JW, et al. Definition and classification of hyperkinetic movements in childhood. *Movement Disorders*. 2010;25(11):1538-49.
10. Sanger TD, Delgado MR, Gaebler-Spira D, Hallett M. Definitions of disorders causing hypertonia in childhood. *Pediatrics*. 2003;111(1):e87-89.
11. Palisano R, Rosenbaum P, Walter S, Russell D, Wood E, Galuppi B. Development and reliability of a system to classify gross motor function in children with cerebral palsy. *Dev Med Child Neurol*. 1997;39(4):214-23.
12. Palisano R, Rosenbaum P, Bartlett D, Livingston MH. *GMFCS- R & E Gross Motor Function Classification System Expanded and Revised, 2007*: CanChild Centre for Childhood Disability, McMaster University, Hamilton, ON; 2007.
13. Palisano RJ, Rosenbaum P, Bartlett D, Livingston MH. Content validity of the expanded and revised Gross Motor Function Classification System. *Dev Med Child Neurol*. 2008;50(10):744-50.
14. Rosenbaum PL, Palisano RJ, Bartlett DJ, Galuppi BE, Russell DJ. Development of the Gross Motor Function Classification System for cerebral palsy. *Dev Med Child Neurol*. 2008;50(4):249-53.
15. Rosenbaum PL, Walter SD, Hanna SE, Palisano RJ, Russell DJ, Raina P, Wood E, Bartlett DJ, Galuppi BE. Prognosis for gross motor function in cerebral palsy: creation of motor development curves. *JAMA*. 2002;288(11):1357-1360.
16. Sackett DL, Straus SE, Richardson WS et al. *Evidence Based Medicine. How to Practice and Teach EBM. Second Edition ed*: Churchill Livingstone; 2000.
17. Brinkmann J. The role of expertise in clinical decision-making: is experience evidence? *O&P Edge*. 2019;24-30.
18. Stuberg W. The aims of lower extremity orthotic treatment in cerebral palsy: a critical review of the literature. In: Condie DN, Meadows CB, editors. *Report of a Consensus Conference on the Lower Limb Orthotic Management of Cerebral Palsy*; Copenhagen: ISPO; 1995. p. 27-34.
19. Morris C. A review of the efficacy of lower-limb orthoses used for cerebral palsy. *Dev Med Child Neurol*. 2002;(44):205-211.
20. Bowers R, Ross K. A review of the effectiveness of lower limb orthoses used in cerebral palsy. In: Morris C, Condie D, editors. *Recent Developments in Healthcare for Cerebral Palsy: Implications and Opportunities for Orthotics*. Copenhagen, Denmark: International Society for Prosthetics and Orthotics; 2009. p. 235-97.
21. Neto HP, Collange Grecco LA, Galli M, Santos Oliveira C. Comparison of articulated and rigid ankle-foot orthoses in children with cerebral palsy: a systematic review. *Pediatr Phys Ther*. 2012;24(4):308-12.
22. Bourseul J-S, Lintanf M, Saliou P, Brochard S, Pons C. Effect of ankle-foot orthoses on gait in children with cerebral palsy: A meta-analysis. *Annals Phys Rehab Med*. 2016;59:E5-E9.
23. Aboutorabi A, Arazpour M, Ahmadi Bani M, Saeedi H, Head JS. Efficacy of ankle foot orthoses types on walking in children with cerebral palsy: A systematic review. *Annals Phys Rehab Med*. 2017;60(6):393-402.
24. Lintanf M, Bourseul J-S, Houx L, Brochard S, Pons C. Effect of ankle-foot orthoses on gait, balance and gross motor function in children with cerebral palsy: a systematic review and meta-analysis. *Clin Rehab*. 2018;32(9):1175-88.
25. Betancourt JP, Eleeh P, Stark S, Jain NB. Impact of ankle-foot orthosis on gait efficiency in ambulatory children with cerebral palsy: a systematic review and meta-analysis. *Am J Phys Med Rehab*. 2019;98(9):759-70.
26. Stevens P, DiBello S. Established indications, benefits and shortcomings of lower-limb orthoses for children with cerebral palsy: a clinical practice guideline. *International J Prosth Orthot*. 2019;31(2, Supp 2, 51.).
27. Owen E. The importance of being earnest about shank and thigh kinematics especially when using ankle-foot orthoses. *Prosth Orthot Int*. 2010;34(3):254-69.
28. World Health Organisation (WHO). *International Classification of Functioning, Disability and Health: ICF Short Version 2001*.

29. Choi H, Wren TAL, Steele KM. Gastrocnemius operating length with ankle foot orthoses in cerebral palsy. *Prosthet Orthot Int*. 2017; 41(3): 274-285.
30. Kerkum YL, Buizer AI, van den Noort JC, Becher JG, Harlaar J, Brehm MA. The effects of varying ankle foot orthosis stiffness on gait in children with spastic cerebral palsy who walk with excessive knee flexion. *PLOS ONE*. 2015; 10(11):e0142878.
31. Kerkum YL, Brehm M-A, van Hutten K, van den Noort JC, Harlaar J, Becher JG, Buizer AI. Acclimatization of the gait pattern to wearing an ankle-foot orthosis in children with spastic cerebral palsy. *Clin Biomech*. 2015; 30(6):617-622.
32. Kerkum YL, Harlaar J, Buizer AI, van den Noort JC, Becher JG, Brehm M-A. An individual approach for optimizing ankle-foot orthoses to improve mobility in children with spastic cerebral palsy walking with excessive knee flexion. *Gait Posture*. 2016;46 :104-111.
33. Danino B, Erel S, Kfir M, Khamis S, Batt R, Hemo Y, Wientroub S, Hayek S. Influence of the orthosis on the foot progression angle in children with spastic cerebral palsy. *Gait Posture*. 2015 Oct; 42(4):518-22.
34. Butler PB, Thompson N., Major RE. Improvement in walking performance of children with cerebral palsy: preliminary results. *Dev Med Child Neurol*. 1992; 34:567-576.
35. Simon S, Deutsch SD, Nuzzo RM, Mansour MJ, Jackson JL, Koskinen M, Rosenthal RK. (1978) Genu recurvatum in spastic cerebral palsy. *J Bone Joint Surg*. 1978; 60-A (7):882-894.
36. International Organization for Standardization (ISO) 13404. *Prosthetics and orthotics – Categorization and description of external orthoses and orthotic components*; 2007.
37. Eddison N, Mulholland M, Chockalingam N. Do research papers provide enough information on design and material used in ankle foot orthoses for children with cerebral palsy? A systematic review. *J Child Ortho*. 2017;11(4):263-71.
38. Owen E. Chapter 21. Segmental Kinematic Approach to Orthotic Management. In: *Physical Therapy for Children With Cerebral Palsy: An Evidence-Based Approach*. 1st Ed. Rahlin ME, editor: SLACK Inc; 2016.
39. Owen E. Call to action: clinical algorithms for the prescription of ankle-foot orthoses are needed: a commentary on “physical therapists’ use of evaluation measures to inform the prescription of ankle-foot orthoses for children with cerebral palsy”. *Phys & Occup Ther In Pediat*. 2019;39(3):254-258.
40. Brehm M, Bus SA, Harlaar J, Nollet F. A candidate core set of outcome measures based on the International Classification of Functioning, Disability and Health for clinical studies on lower limb orthoses. *Prosth & Orthotics Int*. 2011;35(3):269-77.
41. Schiariti V, Selb M, Cieza A, O'Donnell M. International Classification of Functioning, Disability and Health Core Sets for children and youth with cerebral palsy: a consensus meeting. *Dev Med Child Neuro*. 2015;57(2):149-58.
42. Schiariti V, Mahdi S, Bölte S. International Classification of Functioning, Disability and Health Core Sets for cerebral palsy, autism spectrum disorder, and attention-deficit-hyperactivity disorder. *Dev Med Child Neuro*. 2018;60(9):933-41.
43. Owen E. From stable standing to rock and roll walking. part 1. designing, aligning and tuning orthoses for standing stepping and gait. *Association of Paediatric Chartered Physiotherapists Journal*. 2014;5(1):7-18.
44. Owen E. From stable standing to rock and roll walking (part 2) designing, aligning and turning orthoses for standing stepping and gait. *Association of Paediatric Chartered Physiotherapists Journa*. 2014;5(2):4-16.
45. Owen E. Editorial: To tune or not to tune – is that the question? *J Prosthet Orthot*. 2019; 31(3):170-171.
46. Owen E. Editorial. Defining what we do. *J Prosthet Orthot* 2018;30 (1):2-4.
47. Müller S, Carlsohn A, Müller J, Baur H, Mayer F. Static and dynamic foot characteristics in children aged 1–13 years: A cross-sectional study. *Gait Posture*. 2012; 35(3):389-394.
48. Staheli LT. *Fundamentals of Pediatric Orthopedics*, 5th Edition. Philadelphia, USA; Lippincott Williams and Wilkins. 2015.
49. Weber D. (1991) Use of Hinged AFO for Children with Spastic Cerebral Palsy. *J Assoc Child Prosthet Orthot Clin*. 1991; 25:61-65.
50. Robin J, Graham HK, Selber P, Dobson F, Smith K, Baker R. Proximal femoral geometry in cerebral palsy. A population-based cross-sectional study. *J Bone Joint Surg Br*. 2008; 90(10):1372-1379.

©2020 by the APTA Academy of Pediatric Physical Therapy, American Physical Therapy Association, 1020 N Fairfax St, Suite 400, Alexandria, VA 22314, www.pediatricapta.org.

Supported by the Practice Committee of APTA Pediatrics. Developed by expert contributors Elaine Owen, MBE, MSc, SRP, MCSP and Kristie Bjornson, PT, PhD, MS. Illustrations by Elaine Owen and Peter Bull Art Studio, UK. © Elaine Owen. Special thanks to Kathy David, PT, MS for her contributions to the previous version of this fact sheet.

The APTA Academy of Pediatric Physical Therapy provides access to these member-produced fact sheets and resources for informational purposes only. They are not intended to represent the position of APTA Pediatrics or of the American Physical Therapy Association.